

All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Books

Search PubMed

for

Go

Clear

Limits Preview/Index History Clipboard Details

Display Abstract

Show 20

Sort by

Send to

About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

Special Queries

LinkOut

My NCBI (Cubby)

Related Resources

Order Documents

NLM Catalog

NLM Gateway

TOXNET

Consumer Health


Clinical Alerts

ClinicalTrials.gov

PubMed Central

1: Rapid Commun Mass Spectrom. 2005;19(15):2123-30.

Related Articles, Links

**Comparison of negative and positive ion electrospray ionization mass spectra of calmodulin and its complex with trifluoperazine.****Watt SJ, Oakley A, Sheil MM, Beck JL.**Department of Chemistry, University of Wollongong, NSW 2522, Australia.
jbeck@uow.edu.au

The protein calmodulin (apoCaM) undergoes a conformational change when it binds calcium. This structure of the protein (Ca₄CaM) is a dumbbell-shaped molecule that undergoes a further profound conformational change on binding of the antipsychotic drug trifluoperazine (TFP). Experimental conditions were developed to prepare samples of apoCaM, Ca₄CaM and Ca₄CaM/TFP that were substantially free of sodium. The effects of the conformational changes of calmodulin on the charge-state distributions observed in positive ion and negative ion electrospray ionization (ESI) mass spectra were examined. Conversion of apoCaM into Ca₄CaM was concomitant with a change in the negative ion ESI mass spectrum whereby the 16- ion was the most abundant ion observed for the apo form and the 8- ion was the most abundant for the complex. In contrast, in the positive ion ESI mass spectra of apoCaM and Ca₄CaM, the most abundant species in each case was the 8+ ion. When a complex of Ca₄CaM with TFP was prepared, the most abundant species was the 5+ ion. This is consistent with a conformational change of Ca₄CaM that rendered some basic sites inaccessible to ionization in the ESI process. Using the same Ca₄CaM/TFP mixture, no complex with TFP was observed in negative ion ESI mass spectra. These observations are discussed in the context of the structural changes that are known to occur in calmodulin, and suggestions are made to explain the apparently conflicting data. The results reported here reflect on the validity of using differences in charge-state distributions observed in ESI mass spectra to assess conformational changes in proteins. (c) 2005 John Wiley & Sons, Ltd.

PMID: 15988725 [PubMed - in process]

Display Abstract

Show 20

Sort by

Send to